arXiv:1408.2682v2 [math.AG] 15 Jun 2015

Special embeddings of symmetric varieties and their Borel orbits

Mahir Bilen Can[1], Roger Howe[2], and Lex Renner[3]

[1]Tulane University, New Orleans; mcan@tulane.edu

[2]Yale University, New Haven; howe@yale.edu

[3]University of Western Ontario; lex@uwo.edu

August 5, 2014

Abstract

We determine when an antiinvolution on a semisimple linear algebraic group G extends to an antiinvolution on a J-irreducible monoid M whose group of invertible elements is G. Using this information, we study a special class of compactifications of symmetric varieties first considered by Thierry Vust. Generalizing the work of Springer on involutions, we describe the parametrizing sets of Borel orbits in these special embeddings.

Key words: Borel orbits, reductive monoids, symmetric varieties

AMS-MSC: 14M17, 20G05, 20M32

1 Introduction

Through out this paper, $\mathbf{k}$ denotes an algebraically closed field of characteristic zero, and G denotes a linear algebraic group defined over $\mathbf{k}$. We assume, unless explicitly stated otherwise, that G is connected.

A closed subgroup $H \subseteq G$ is called spherical if there exists a Borel subgroup $B \subseteq G$ such that BH is an open subset of G. Equivalently, the coset space G/H on which G-acts algebraically from the left has a dense B-orbit. More generally, a normal G-variety X is called spherical (or, more descriptively, *G-spherical*) if B has a dense orbit in X. By the works of Brion [1], Popov [20], and Vinberg [36], it is known that

X is spherical if and only if X is comprised of finitely many B-orbits. Such varieties are abundant and useful in many branches of mathematics. Toric varieties as well as homogeneous varieties are among their well known and widely used examples. They arise naturally in the study of real forms of linear groups. It is shown by Matsuki in [17] (when $\mathbf{k}$ is the field of complex numbers) and by Springer in [33] (for general $\mathbf{k}$) that the fixed subgroup

$$G^\theta = \{g \in G : \ \theta(g) = g\}$$

of an automorphism $\theta : G \to G$ such that $\theta^2 = id$ is a spherical subgroup. We refer to such an automorphism as an *involution*, even though there is a low risk of confusing it with the 'involution in a Weyl group'. Nevertheless, the understanding of which one is meant will be clear from the context.

A coset variety of the form G/G^θ is called a symmetric variety. Over the field of real numbers, such manifolds are known as "Riemannian symmetric spaces." The isomorphism types of compact Riemennian spaces are described by Cartan in [5], and more uniformly by Huang and Leung in [13]. Further prominent examples of spherical varieties appear in the modern theory of group compactifications: A variety X, which is assumed to be normal, is called an embedding of G/H if it has an open subset that is isomorphic to G/H. Clearly, in this case, if H is a spherical subgroup, then X is a spherical variety. If G is a product group of the form $K \times K$ and H is the diagonal copy of K in G, then G/H is isomorphic to K as a $K \times K$-variety. In [30], Rittatore shows that an affine $K \times K$-variety X is an embedding of the group K if and only if X is a linear algebraic monoid, that is to say a closed, unital subsemigroup of the monoid of $n \times n$ matrices for some n. Moreover, in this case, K is the group of invertible elements of the monoid X.

The overarching goal of our paper is to gain a deeper understanding of the spherical varieties. Roughly speaking our main results are; a construction of a "new" class of spherical embeddings of symmetric varieties using linear monoids, and a description of the sets parameterizing Borel orbits therein. Before getting into the details of our results, we present an observation, along the way of which we introduce some concepts that we need in the sequel:

Suppose G is semisimple and of adjoint type, and $\theta : G \to G$ is an involutory automorphism. Let $\widetilde{H}$ denote the normalizer of the fixed subgroup $H = G^\theta$ in G. According to [7], there exists a unique normal G-variety X such that

i) there exists an open G-orbit $X_0 \subset X$ that is isomorphic to $G/\widetilde{H}$,

ii) the complement $X - X_0$ is a finite union $\cup_{\alpha \in S} X_\alpha$ of smooth, G-stable boundary divisors with normal crossings such that $X_\alpha \cap X_\beta \neq \emptyset$ for all $\alpha, \beta \in S$,

iii) any G-orbit closure is of the form $\cap_{\alpha \in I} X_\alpha$ for some subset $I \subset S$, where S is as in ii).

The variety X is called the wonderful compactification of $G/\widetilde{H}$. In the case when G is the doubled group $K \times K$, and the involution θ is $\theta((h,k)) = (k,h)$ for $(h,k) \in K \times K$, the wonderful compactification of $G/\widetilde{H}$ is closely related to a particular linear algebraic monoid. To explain we recall some standard algebraic monoid definitions. A *reductive* monoid is, by definition, a linear algebraic monoid with a reductive group of invertible elements. A *J-irreducible* monoid is a reductive monoid M with a unique closed $G \times G$-orbit that has at least two elements including the zero element. Such monoids are abundant in representation theory. Let (ρ, V) be a finite dimensional irreducible representation of a semisimple group G_0. The image of ρ_0 lies in $\mathrm{End}(V)$, the affine space of linear operators on V. By abusing notation, we denote by $\mathbf{k}^*$ the group of nonzero scaling operators on V and let G denote the product subgroup $\mathbf{k}^* \cdot \rho(G_0)$ of $\mathrm{GL}(V)$. The Zariski closure $\overline{G} \subseteq \mathrm{End}(V)$ is a J-irreducible monoid. To obtain the wonderful compactification of G_0, a semisimple group of adjoint type, we start with a representation V with its highest weight in general position such that $\dim(V \otimes V)^{G_0} = 1$. Normalizing the projectivization of the associated J-irreducible monoid gives us the wonderful compactification of G_0. See [26].

In this paper we practice the following yoga: interesting compactifications come from semigroup theory, and (somewhat) conversely, interesting monoids arise as the embeddings of groups. To explain our mantra, we start with embedding our symmetric variety G/H into a J-irreducible monoid.

Let $\tau : G \to G$ be the morphism defined by

$$\tau(g) = g\theta(g)^{-1} \tag{1.1}$$

It is shown by Vust in [37] and later simplified by Richardson in [28] that the image of τ in G, denoted by $P = \{\tau(g) : \ g \in G\}$, is isomorphic to G/H. Observe that if G is equal to $K \times K$ and the involution on G is defined by $\theta(h,k) = (k,h)$ for all $k, h \in K$, then P is equal to the whole variety K.

Next, we consider a J-irreducible monoid M whose group of invertible elements is G. Clearly, the closure of P in M gives us an embedding of G/H. At this point it is desirable to determine if the defining involution of P has some sort of extension to M, or not. Paraphrasing this as a question, we ask:

"For which monoids M do we have an involution to work with?"

Our first main result answers this question in the absence of invertibility.

Theorem 1. Let θ be an involutory automorphism of G and let H be the fixed subgroup $H = G^\theta$. Let (ρ, V) be an irreducible representation of G with highest weight μ and let M denote the associated J-irreducible monoid. If the induced map θ^* on weights satisfies $\theta^*(\mu) = -\mu$, then there exists a unique morphism $\theta_{an} : M \to M$ such that

1. $\theta_{an}(xy) = \theta_{an}(y)\theta_{an}(x)$ for all $x, y \in M$,

2. θ_{an}^2 is the identity map on M,

3. $\theta_{an}(g) = \theta(g)^{-1}$ for all $g \in G$.

As in the case of homogeneous varieties, closures of orbits of a Borel subgroup in a symmetric variety dictates the underlying geometry (see [2] and [**?**]). In [33], Springer determines the structure of these orbits. The inclusion posets of orbit closures are investigated by Matsuki in [18] (for $\mathbb{C}$), and by Richardson and Springer in [29] (for arbitrary $\mathbf{k}$ with characteristic zero). These results are sharpened and extended over the fields of arbitrary characteristic (not two) by Helminck and Wang in [11], and by Helminck in [10]. Much of what we describe next provides us with a new ground of applications of these results. Before stating our next theorem in this direction, let us briefly summarize what is known for B-orbits in an arbitrary symmetric variety.

Let $T \subseteq B$ be a θ-stable maximal torus of a Borel subgroup B of G. The *Weyl group* W of G is the finite group $N_G(T)/T$, where $N_G(T)$ is the normalizer of T in G. Let $\mathcal{V}$ denote the set of all $g \in G$ such that $\tau(g) \in N_G(T)$, where τ is as in (1.1). It is easy to verify that $\mathcal{V} \subset G$ is closed under the action

$$(t, h) \cdot g = tgh^{-1} \quad \text{for all } t \in T, h \in H, \ g \in G.$$

Let V denote the set of $T \times H$-orbits in $\mathcal{V}$. For $v \in V$, let $x(v) \in \mathcal{V}$ denote a representative of the orbit v. The inclusion $\mathcal{V} \hookrightarrow G$ induces a bijection from V onto the set of $B \times H$-orbits in G. In particular, G is the disjoint union of the double cosets $Bx(v)H$, $v \in V$. See [10].

Suppose now that M is a reductive monoid with an antiinvolution $\theta_{an} : M \to M$. Obviously, θ_{an} composed with the inverting morphism $\iota(g) = g^{-1}$ on G is an involution, denoted by θ, on the unit group G. Let H and P denote, as before, the fixed subgroup $H := G^\theta$ and the closed subvariety of G which is isomorphic to G/H. We define the *twisted action* of G on M by

$$g * m = gm\theta_{an}(g) = gm\theta(g)^{-1}, \qquad g \in G, m \in M. \tag{1.2}$$

P is stable under the action (1.2). In fact, $P = G*1_G$, where 1_G is the identity element of G. Since P is open in its closure $\overline{P}$ in M, $\overline{P}$ is closed under (1.2). In particular, $\overline{P}$ is a spherical variety. An important auxiliary variety for studying symmetric varieties is the subvariety

$$Q := \{g \in G : \ \theta(g) = g^{-1}\}.$$

The following facts about Q are easy to check

1. Q is closed in G;
2. Q is stable under the twisted action (1.2);
3. Q is the fixed subvariety of the antiinvolution θ_{an} on G;
4. $P \subseteq Q$.

Moreover, we know from [33] that Q has only finitely many $B*$-orbits.

Generalizing Q and some related objects enables us to lift many results of Springer from [33] to the monoid setting. First, we observe that Vust's morphism (1.1) has a natural extension to M:

$$\tau : M \longrightarrow M$$
$$\tau(x) = x\theta_{an}(x)$$

Define

$$M_Q := \{m \in M : \ \theta_{an}(m) = m\}. \tag{1.3}$$

Since θ_{an} is a morphism and M is separated, M_Q is a closed subset in M. It follows from the inclusions $P \subseteq Q \subseteq M_Q$ that $\overline{P} \subseteq \overline{Q} \subseteq M_Q$.

G acts on the sets $P, Q, \overline{P}, \overline{Q}$, and on M_Q by the same formula $g * m = gm\theta_{an}(g)$. Our goal is to give a satisfactory description of the $B*$-orbits in these varieties. Note that, $B*$-orbits in G are in one-to-one correspondence with the B-orbits in G/H.

Theorem 2. Let $T \subseteq G$ be a θ-stable maximal torus in a θ-stable Borel subgroup $B \subset G$, and let $\overline{N}$ denote the Zariski closure in M of the normalizer $N_G(T)$. The following sets are finite and they are in bijection with each other:

1. $B*$-orbits in $\overline{Q}$ (respectively, $B*$-orbits in $\overline{P}$),
2. $T \times H$-orbits in $\tau^{-1}(\overline{N} \cap \overline{Q})$ (respectively, $T \times H$-orbits in $\tau^{-1}(\overline{N} \cap \overline{P})$).

Before closing our introduction by presenting the contents of our paper, we comment on the consequences of Theorem 2. The *Renner monoid*, $R = \overline{N}/T$ generalizes the notion of Weyl group from linear algebraic groups to reductive monoids [25]. Its elements are in bijection with the $B \times B$-orbits in M, and its system of idempotents encodes crucial information on the structure of a suitable torus embedding. Its inverse semigroup structure encapsulates many interesting combinatorial properties. See [16]. The partial ordering on R that is induced by the inclusion relations among the closures of $B \times B$-orbits plays an important role for the geometry of M and for the related algebras. See [31], [22], [23], and [19]. Theorem 2 opens a way for many

related enumerative and combinatorial questions. As consequence of Theorem 2 we know that a certain subset of R can be used to study $B*$-orbits in $\overline{P}$. In fact, in special cases this subset parametrizes the full set of $B*$-orbits (see the examples in Section 5). There are many open questions here. For example, is there a formula for the size of this subset of R? Of course, the same question makes sense for the $B*$-orbits in $\overline{Q}$.

We structured our paper as follows. In Section 2, we recall some well-known facts from the representation theory of semisimple linear groups, in particular, the theory of weights. Also in the same section, we review the basics of the theory of linear algebraic monoids and review the convex geometry of a maximal tori in the monoid. In Section 3, after recalling the theory of root system related to symmetric varieties, we prove our first main result, Theorem 1. The definition of a special compactification and its basic properties are given in Section 4. In Section 5, we obtain the characterization of the parametrizing sets of Borel orbits in the embeddings that are mentioned earlier. In particular we prove our Theorem 2 as a corollary of the related result which describes the set of Borel orbits in M_Q. We close our paper with some final remarks in Section 6.

Acknowledgement. The authors thank the anonymous referee for her/his comments and suggestions which improved the quality of the paper drastically. The first author is partially supported by the Louisiana Board of Regents Research and Development Grant 549941C1 and by the N.S.A. Grant H98230-14-1-0142.

2 Preliminaries

In this section, we introduce our basic notation and give the necessary background for algebraic groups and monoids. For representation theory related notation, we follow the books [14, 15], and for monoids we follow [27] and [21].

We denote by GL_n the general linear group of invertible $n\times n$ matrices, and denote by Mat_n the monoid of $n \times n$ matrices. A linear algebraic group is a closed subgroup of GL_n. More generally, a linear algebraic monoid is a closed submonoid of Mat_n. We always assume that a linear algebraic group is connected.

The Lie algebra of a linear algebraic group G is denoted by $\mathrm{Lie}(G)$, and the *adjoint representation* $(\rho, V) = (\mathrm{Ad}, \mathrm{Lie}(G))$ of G is defined by the action

$$g \cdot A = gAg^{-1} \quad \text{for all } g \in G, A \in \mathrm{Lie}(G).$$

G is called simple if it has no non-trivial, normal, connected subgroup. Equivalently, the Lie algebra of G has no non-trivial abelian ideals. In this case, $\mathrm{Lie}(G)$ is said to be a simple Lie algebra. G is called semisimple, if its Lie algebra is *semisimple*, that is

to say Lie(G) is a sum of simple Lie algebras. More generally, G is called reductive if its Lie algebra is the sum of an abelian and a semisimple Lie algebra. Throughout the paper, we consider reductive groups, only. Our final general assumption on groups and monoids is that all representations are rational and finite dimensional.

2.1 Weights

If $T \subseteq G$ is a maximal torus, then a *weight* of T on a representation (π, V) of G is a character $\lambda : T \to \mathbf{k}^*$ such that the subspace $V_\lambda = \{v \in V : \ \pi(t) \cdot v = \lambda(t)v \text{ for all } t \in T\}$ is nonempty. We denote by $X(V)$ the (finite) set of weights of a given representation (π, V). Collection of all weights (from all representations) of T forms a free abelian group, denoted by $X(T)$. The real vector space $X(T) \otimes_{\mathbb{Z}} \mathbb{R}$ is denoted by E. For a Borel subgroup B containing T, we have the following additional notation:

$$\begin{aligned} \Phi &\subset X(T) \text{ the set of weights of the adjoint representation,} \\ \Delta &\subset \Phi \text{ the set of simple roots determined by } (B, T), \\ \Phi^+ &\subset \Phi \text{ the subset of positive roots associated with } \Delta. \end{aligned}$$

Recall that the Weyl group of a semisimple group (more generally, of a reductive group) G is the finite quotient group $W = N_G(T)/T$, where $N_G(T)$ is the normalizer of a maximal torus T in G. There is a faithful representation of W on E by which W is recognized as a "reflection group." In other words, there exists a W-invariant inner product $(\ ,\ ) :\ E \times E \to \mathbb{R}$ and the group generated by the orthogonal transformations

$$\sigma_\alpha(\mu) = \mu - \frac{2(\mu, \alpha)}{(\alpha, \alpha)}\alpha, \qquad \text{for all } \mu \in E, \alpha \in \Phi \tag{2.1}$$

is isomorphic to W.

Let $\langle\ ,\rangle : E \times E \to \mathbb{R}$ denote the pairing defined by $\langle \lambda, \mu \rangle = 2(\lambda, \mu)/(\mu, \mu)$, $\mu, \lambda \in E$. An *abstract weight* in E is a vector $\lambda \in E$ such that $\langle \lambda, \alpha \rangle \in \mathbb{Z}$ for all $\alpha \in \Phi$. In particular $X(T)$ consists of abstract weights. The set of all abstract weights in E forms a lattice (of full rank) denoted by Λ, and G is called *simply-connected*, if $\Lambda = X(T)$. The sublattice of $X(T)$ generated by the set of all roots is called the root lattice and it is denoted by Λ_r. G is called of adjoint type if $X(T) = \Lambda_r$.

If Δ is $\{\alpha_1, \ldots, \alpha_\ell\}$, then the *fundamental dominant weights* ω_i's are defined by the requirements $\langle \omega_i, \alpha_j \rangle = \delta_{i,j}$ for $i, j = 1, \ldots, \ell$. They form a basis for the lattice Λ, as well as for the vector space E. A vector $\lambda \in E$ is called dominant if all c_i's in its expansion $\lambda = \sum_{i=1}^{\ell} c_i \omega_i$ are nonnegative integers. If for all $i = 1, \ldots, \ell$ the integer c_i is positive, then λ is called a regular (dominant) weight. By its representation defined

in (2.1), W acts on Λ, furthermore, each $\lambda \in \Lambda$ is W-conjugate to a unique dominant weight.

There is an important partial ordering $\leqslant$ on E defined by $\mu \leqslant \lambda$ if $\lambda - \mu$ is a nonnegative integer combination of positive roots. W permutes the weights $X(V)$ of (π, V), and there is a unique weight $\lambda \in X(V)$ such that $\mu \leqslant \lambda$ for all $\mu \in X(V)$. In this case, λ is called the highest weight of the representation (π, V).

2.2 Reductive monoids

The purpose of this section is to introduce the notation of the reductive monoids. We focus mainly on the orbit structure of these monoids. In addition to Renner and Putcha's books mentioned above, we recommend more recent semi-expository article by Brion [3]. For an introduction from a combinatorial point of view with many explicit examples we suggest the survey by Solomon [32]. For Renner monoids, we recommend Cao, Li and Li's survey article [16].

Let M be a linear algebraic monoid. The set $G = G(M)$ of invertible elements of M is an algebraic group. If G is a reductive group and M is an irreducible variety, then M is said to be a reductive monoid. In a reductive monoid, the data of the Weyl group W of (G, T) and the set $E(\overline{T})$ of idempotents of the embedding $\overline{T} \hookrightarrow M$ combine to become a finite inverse semigroup $R = \overline{N_G(T)}/T \cong W \cdot E(\overline{T})$ with unit group W and idempotent set $E(R) = E(\overline{T})$. (Here, $E(\cdot)$ stands for the idempotent set of a semigroup.)

The*Bruhat-Chevalley order* on W is defined by

$$x \leq y \quad \text{if and only if} \quad BxB \subseteq \overline{ByB}.$$

Similarly, on the Renner monoid R of M, the Bruhat-Chevalley order is defined by

$$\sigma \leq \tau \quad \text{if and only if} \quad B\sigma B \subseteq \overline{B\tau B}. \tag{2.2}$$

The induced poset structure on W, which is induced from R is the same as the original Bruhat poset structure on W.

There is a canonical partial order $\leq$ on the set of idempotents $E(M)$ of M (hence on $E(\overline{T})$) defined by

$$e \leq f \quad \text{if and only if} \quad ef = e = fe. \tag{2.3}$$

Notice that $E(\overline{T})$ is invariant under the conjugation action of the Weyl group W. A subset $\Lambda \subseteq E(\overline{T})$ is called a cross-section lattice (or, a *Putcha lattice*) if Λ is a set of representatives for the W-orbits on $E(\overline{T})$ and the bijection $\Lambda \to G\backslash M/G$ defined

by $e \mapsto GeG$ is order preserving. There is a close relationship between cross-section lattices and Borel subgroups.

The *right centralizer of* Λ *in* G, denoted by $C_G^r(\Lambda)$, is the subgroup

$$C_G^r(\Lambda) = \{g \in G : \ ge = ege \text{ for all } g \in G\}.$$

It turns out that given a cross-section lattice Λ, there exists a unique Borel subgroup $B = B(\Lambda)$ such that $B = C_G^r(\Lambda)$. Conversely, any cross-section lattice is of the form $\Lambda = \Lambda(B) = \{e \in E(\overline{T}) : \ Be = eBe\}$ for a unique Borel subgroup B containing T.

The decomposition $M = \bigsqcup_{e \in \Lambda} GeG$ into $G \times G$ orbits has a counterpart on the Renner monoid. Namely, the finite monoid R can be written as a disjoint union

$$R = \bigsqcup_{e \in \Lambda} WeW$$

of $W \times W$ orbits, parametrized by the cross section lattice. Moreover, the partial order (2.3) on Λ agrees with the order induced from Bruhat-Chevalley order (2.2).

$E(\overline{T})$ is a relatively complemented lattice, anti-isomorphic to a face lattice of a convex polytope. For $\overline{T}$ contained in a J-irreducible monoid, the associated polytope is described explicitly in Section 2.3. Let Λ be a cross section lattice in $E(\overline{T})$. The Weyl group of T (relative to $B = C_G^r(\Lambda)$) acts on $E(\overline{T})$, and furthermore

$$E(\overline{T}) = \bigsqcup_{w \in W} w\Lambda w^{-1}.$$

Let S be a semigroup and let $M = S^1$ be the monoid obtained from S by adding a unit element if it is not already present. Let $a, b \in M$. The following are four of the five equivalence relations which are collectively known as Green's relations. They are of utmost importance for semigroup theory.

1. $a \ \mathscr{L} \ b$ if $Ma = Mb$.
2. $a \ \mathscr{R} \ b$ if $aM = bM$.
3. $a \ \mathscr{J} \ b$ if $MaM = MbM$.
4. $a \ \mathscr{H} \ b$ if $a \ \mathscr{L} \ b$ and $a \ \mathscr{R} \ b$.

It turns out the unit group G of a reductive monoid M is big in the sense that $a \ \mathscr{L} \ b$ if $Ga = Gb$, $a \ \mathscr{R} \ b$ if $aG = bG$, and $a \ \mathscr{J} \ b$ if $GaG = GbG$. Furthermore, a cross section lattice is a representative for the set of $\mathscr{J}$-classes in M (see Proposition 6.1 of [21]). A reductive monoid M is called J-irreducible if M has a unique, nonzero, minimal $G \times G$-orbit.

Among the submonoids that play a prominent role for understanding the structure of M are those of the form eMe, $e \in \Lambda$. It is turns out eMe is a reductive monoid itself, and moreover, its group of invertible elements, H_e is equal to $eC_G(e)$. For more on special submonoids, see Chapter 7 of [21].

2.3 J-irreducible monoids and their cones

Let G_0 be a simple linear algebraic group and let T_0 be a maximal torus in G_0. If (ρ_0, V) is an n-dimensional irreducible representation of G_0 such that $\dim \ker(\rho_0) = 0$, then the group $\mathbf{k}^* \cdot \rho_0(G_0)$, which we denote by G is reductive. Up to isomorphism T_0 and $\rho_0(T_0)$ differ by a finite set of central elements, so, when there is no danger of confusion, by abusing notation, we denote the image $\rho_0(T_0)$ by T_0, also. Let $T \subseteq G$ be a maximal torus containing T_0, and let $\mathbf{T} \subset \mathrm{GL}(V)$ denote an n-dimensional maximal torus containing T. Accordingly, we have a nested sequence of Euclidean spaces:

$$E_0 = X(T_0) \otimes_{\mathbb{Z}} \mathbb{R} \subset E = X(T) \otimes_{\mathbb{Z}} \mathbb{R} \subset \mathbf{E} = X(\mathbf{T}) \otimes_{\mathbb{Z}} \mathbb{R}.$$

Note that $\dim E = \dim E_0 + 1$.

If ε_i (for $i = 1, \ldots, n$) denotes the standard i-th coordinate function on $\mathbf{T}$, then $\{\varepsilon_1, \ldots, \varepsilon_n\}$ is a basis for $\mathbf{E}$, and E is spanned by the restrictions $\varepsilon_i|_T$, $i = 1, \ldots, n$. Let $\chi \in X(T)$ denote the restriction of the character whose n-th power is the determinant on $\mathrm{GL}(V)$. Stated differently in additive notation, χ is the restriction to T of the rational character $\frac{1}{n}(\varepsilon_1 + \cdots + \varepsilon_n)$. We denote $\varepsilon_i|_T$ by χ_i and set

$$\widetilde{\chi}_i := \chi_i - \chi \quad \text{for } i = 1, \ldots, n.$$

If K is an arbitrary group, then the center of K is customarily denoted by $Z(K)$. In our case, since $Z(G) = \mathbf{k}^* \cdot Z(G_0)$, the character group of $Z(G)$ is generated by one element, which is χ. Thus, $E = \mathbb{R}\chi \oplus E_0$. In fact, χ vanishes on T_0. It follows from these observations that

1. $\{\widetilde{\chi}_1, \ldots, \widetilde{\chi}_{n-1}, \chi\}$ spans E;
2. if $x \in T$ lies in $T_0 \subset T$, then $\widetilde{\chi}_i|_{T_0}(x) = \chi_i(x)$;
3. $\{\widetilde{\chi}_1, \ldots, \widetilde{\chi}_{n-1}\}$ spans E_0.

In this paper, we are interested in the J-irreducible monoids that come from a faithful representation.

Definition 3. Let (ρ_0, V) be an irreducible representation of G_0. The *J-irreducible monoid associated with* (ρ_0, V) is the affine variety $\overline{\mathbf{k}^* \cdot \rho_0(G_0)}$ together with its monoid structure induced from $\mathrm{End}(V)$.

Remark 4. The definition of a J-irreducible monoid which is given at the end of Section 2.2 agrees with Definition 3, see Lemma 7.8 of [27].

The representation (ρ_0, V) of G_0 gives a family of representations for G: For each natural number $k \in \mathbb{N}$, there is a representation defined by the action

$$zg \cdot v = z^k(\rho_0(g)(v)) \in V, \tag{2.4}$$

where $z \in \mathbf{k}^*$, $g \in G_0$, and $v \in V$. Another such simple but useful observation is that, since $G_0 \overset{\rho_0}{\cong} (G, G)$, the Weyl group W of (G_0, T_0) is isomorphic to that of the pair (G, T).

Lemma 5. [Proposition 3.5 in [24]] Let (ρ_0, V) be an irreducible representation of G_0. If λ is the T_0-highest weight of (ρ_0, V) and $\mathcal{P}$ denotes the convex hull of $\{w \cdot (\chi + \lambda) : w \in W\} = \{\chi + w \cdot \lambda : \ w \in W\}$, then the set of weights of T (with respect to (2.4) with $k = 1$, for simplicity) is contained in $\mathcal{P}$.

A *strongly convex rational polyhedral cone* in $\mathbb{R}^m$ (for some $m > 0$) is a cone c whose apex is at the origin and such that

- c is generated by finitely many vectors which belong to a sublattice of $\mathbb{R}^m$ ("rationality");
- c does not contain a line passing through the origin ("strong convexity").

Now, if σ denotes the strongly convex rational polyhedral cone generated by the vectors $\chi_1, \ldots, \chi_n$ in E, then it follows from the discussion after Lemma 3.1 and the proof of Proposition 3.5 in [24] that σ is equal to the cone $\sum_{w \in W} \mathbb{R}^+ w \cdot \lambda$. In other words, σ is a strongly convex cone over $\mathcal{P}$. (Figure 4 in page 202 of [24] is a good depiction of what is meant here.)

Example 6. Let G_0 stand for SL_n, the special linear group consisting of $n \times n$ matrices with determinant 1, and let T_0 denote the maximal torus of diagonal matrices in G_0. The *first fundamental representation* $\rho_0 : G_0 \to \mathrm{GL}(\mathbf{k}^n) \simeq \mathrm{GL}_n$ is defined by

$$\rho_0(x) = x, \quad \text{for all} \quad x \in G_0.$$

Clearly, $G = \mathbf{k}^* \cdot \rho_0(G_0)$ is equal to $\mathrm{GL}(\mathbf{k}^n)$, and therefore, $T = \mathbf{T}$. It is straightforward

to verify in this case that $\chi_i = \varepsilon_i$ and that the weight ω_1[1] defined by

$$\omega_1 := \chi_1 - \frac{1}{n}(\chi_1 + \cdots + \chi_n) \tag{2.5}$$

is the T_0-highest weight of $(\rho_0, \mathbf{k}^n)$. Note that

- $\{\chi_1 - \chi, \ldots, \chi_{n-1} - \chi, \chi\}$ is a basis for $X(T)$;
- $\{\widetilde{\chi}_i = \chi_i - \chi : \ i = 1, \ldots, n-1\}$ is a basis for $X(T_0)$.

For $n \in \mathbb{N}$, we use the notation $[n]$ to denote the set $\{1, \ldots, n\}$. The symmetric group of permutations on $[n]$, denoted by S_n, is the Weyl group of both (GL_n, T) and (SL_n, T_0). It acts on χ_i's by permuting their indices;

$$\pi \cdot \chi_i = \chi_{\pi^{-1}(i)} \ \text{ for all } \ \pi \in S_n, \ \text{ and } \ i = 1, \ldots, n. \tag{2.6}$$

We re-write the n-th coordinate function in the following way:

$$\begin{aligned} \chi_n &= n\chi - (\chi_1 + \cdots + \chi_{n-1}) \\ &= (\chi - \chi_1) + \cdots + (\chi - \chi_{n-1}) + \chi. \end{aligned}$$

Thus, $\widetilde{\chi}_n = \chi_n - \chi = -(\widetilde{\chi}_1 + \cdots + \widetilde{\chi}_{n-1})$. Now it is easily seen that S_n stabilizes E_0, the $\mathbb{R}$-span of $\{\widetilde{\chi}_i : \ i = 1, \ldots, n-1\}$. Let σ denote the cone

$$\sigma = \mathbb{R}^+\chi_1 \oplus \cdots \oplus \mathbb{R}^+\chi_{n-1} \oplus \mathbb{R}^+\chi_n \ \text{ in } \ E \simeq \mathbb{R}^n.$$

Notice that ω_1 of (2.5) is equal to $\widetilde{\chi}_1$, and therefore, $\chi + \omega_1 = \chi_1$. Thus, by (2.6) the convex hull of vectors $w \cdot (\chi + \omega_1)$, $w \in S_n$ is equal to the convex hull of χ_i's, and σ is a strongly convex cone over this polytope. We identify χ_i with the n-tuple $e_i = (0, \ldots, 1, \ldots, 0)$ having a single 1 at the i-th entry, and 0's elsewhere. The usual permutation action of S_n on e_i's agrees with the action of W on χ_i's, hence, we see that $\mathcal{P}$ is the regular $(n-2)$-simplex. See Figure 2.1 for $n = 5$, for which we use the projection from $\mathbb{R}^4$ to $\mathbb{R}^3$.

[1] More generally, for $i = 1, \ldots, n-1$ define

$$\omega_i = \chi_1 + \cdots + \chi_i - \frac{i}{n}(\chi_1 + \cdots + \chi_n).$$

It is a classical result by Cartan that ω_i is the highest weight of the i-th exterior power $\bigwedge^i \mathbf{k}^n$ of the first fundamental representation of SL_n. By analogy, $\bigwedge^i \mathbf{k}^n$ is called the i-th fundamental representation.



Figure 2.1: The polytope of the defining representation $\mathrm{SL}_5 \to \mathrm{GL}(\mathbf{k}^5)$.

Example 7. Observe that $Z(\mathrm{SL}_n)$ is comprised of n-roots of 1. Let G_0 denote the adjoint group $\mathrm{SL}_n/Z(\mathrm{SL}_n)$ and (ρ_0, V) denote its adjoint representation. In other words, V is equal to $\mathfrak{sl}_n$, the space of traceless $n \times n$ matrices, and

$$\rho_0(g)(A) = gAg^{-1} \text{ for all } g \in G_0, \text{ and } A \in V.$$

Since $\ker(\rho_0) \subseteq \mathrm{SL}_n \cap \mathrm{Z}(\mathrm{Mat}_n)$, ρ_0 is faithful. As before, let G denote the extended group $\mathbf{k}^* \cdot \rho_0(G_0) \subset \mathrm{GL}(V)$, and let T denote the maximal torus of G containing T_0. Here, $T_0 \subset G_0$ is the (image of the) maximal torus consisting of diagonal matrices. Note that $G \simeq \mathrm{GL}_n$.

Let $[n]^2$ denote the set $[n] \times [n]$, and let $([n]^2)^*$ denote the set $[n]^2 - \{(n,n)\}$. A canonical Lie algebra basis for $V = \mathfrak{sl}_n$ is given by the ordered set $B = \{F_{i,j} : (i,j) \in ([n]^2)^*\}$ (ordered with respect to lexicographic ordering on the indices), where

$$F_{i,j} = \begin{cases} E_{i,j} & \text{if } 1 \leq i \neq j \leq n, \\ E_{i,i} - E_{i+1,i+1} & \text{if } 1 \leq i = j \leq n-1. \end{cases}$$

Here $E_{i,j}$ is the elementary matrix with 1 at its i,j-th entry, and 0's everywhere else. We identify $\mathrm{GL}(V)$ with GL_{n^2-1} via the basis B. Let $\mathbf{T}$ denote the maximal torus of diagonal matrices in GL_{n^2-1}, and let $\varepsilon_{i,j}$ denote the coordinate function defined by

$$\varepsilon_{i,j}(\mathrm{diag}(a_1, \ldots, a_{n^2-1})) = a_{(n-1)i+j}.$$

The character group $X(\mathbf{T})$ of $\mathbf{T}$ is generated by $\varepsilon_{i,j}$'s. Let $\chi_{i,j}$ denote the restriction of $\varepsilon_{i,j}$ to T. It is a straightforward calculation to show that a generic element y from $T \subset \mathbf{T}$ is of the form $y = cx$, where

$$x = \mathrm{diag}(t_1 t_1^{-1}, t_1 t_2^{-1}, \ldots, t_1 t_n^{-1}, \ldots, t_{n-1} t_1^{-1}, \ldots, t_{n-1} t_n^{-1}, t_n t_1^{-1}, \ldots, t_n t_{n-1}^{-1}) \qquad (2.7)$$

for some diagonal matrix $\mathrm{diag}(t_1, \ldots, t_n) \in \mathrm{SL}_n$, and for some $c \in \mathbf{k}^*$. In particular, $x \in \rho_0(T_0)$. Therefore, $\chi_{i,j}(y) = ct_it_j^{-1}$. Let χ denote the restriction to T of the (n^2-1)-th root of the product of all $\varepsilon_{i,j}$'s:

$$\chi = \left(\prod_{(i,j)\in([n]^2)^*} \chi_{i,j} \right)^{\frac{1}{n^2-1}}.$$

Thus, $\chi(y) = \chi(cx) = c = \det y$ for $y = cx \in T$, where x is as in (2.7).

For $(i,j) \in ([n]^2)^*$, we put $\widetilde{\chi}_{i,j} := \chi_{i,j} - \chi$. We know that the character group $X(T)$ is amply generated by the union $\{\widetilde{\chi}_{i,j} : 1 \le i \ne j \le n\} \cup \{\chi\}$, and that the set $\{\widetilde{\chi}_{i,j} : 1 \le i \ne j \le n\}$ generates $X(T_0)$. Both of $X(T_0)$ and $X(T)$ are sublattices of $X(\mathbf{T}) = \bigoplus_{(i,j)\in([n]^2)^*} \mathbb{Z}\varepsilon_{i,j}$.

The Weyl group $W \simeq S_n$ of (G,T) (hence of (G_0,T_0)) acts on $\widetilde{\chi}_{i,j}$'s by permuting the indices as follows:

$$w \cdot \widetilde{\chi}_{i,j} = \widetilde{\chi}_{w^{-1}i,w^{-1}j} \quad \text{for all } w \in S_n \text{ and } 1 \le i \ne j \le n.$$

Clearly, the determinant is intact by this action. Similar to the case of the previous example, we identify $\widetilde{\chi}_{i,j}$ with a suitable vector in $\mathbb{R}^n$ with entries from $\{-1,0,1\}$:

$$\widetilde{\chi}_{i,j} \leftrightsquigarrow \begin{cases} (0,\ldots,0,1,0\ldots,0,-1,0\ldots0) & \text{if } i<j; \\ (0,\ldots,0,-1,0\ldots,0,1,0\ldots0) & \text{if } i>j. \end{cases} \tag{2.8}$$

Here, the nonzero entries occur only at the i-th and the j-th positions. Note that there exactly $n(n-1)$ such vectors. The identification (2.8) is equivariant in the sense that the action of W on $\widetilde{\chi}_{i,j}$'s agrees with the permutation action of S_n on the entries of the corresponding vectors.

It is well known that the highest weight λ of the adjoint representation of SL_n is equal to $\omega_1 + \omega_{n-1} = \varepsilon_1 - \varepsilon_n$, where ε_i is the i-th coordinate function on T_0. Equivalently,

$$\lambda = \omega_1 + \omega_{n-1} = \widetilde{\chi}_{1,n} = \chi_{1,n} - \chi \in X(T_0) \subset X(\mathbf{T}).$$

Therefore, it follows from (2.8) that the polytope $\mathcal{P}$ (the convex hull of $W \cdot (\lambda + \chi)$) is equal to the convex hull in $\mathbb{R}^n$ of all $0/1$ vectors with exactly two nonzero entries. For example, when $n = 4$ we see, after projecting from $\mathbb{R}^4$ to $\mathbb{R}^3$ that $\mathcal{P}$ is equal to the cube-octahedron depicted in Figure 2.2.

3 Special weights; a proof of Theorem 1

Let G denote a reductive group and θ be an involution on G, Then there exists a θ-stable maximal torus T in G. Moreover, there exists a θ-stable Borel subgroup



Figure 2.2: The polytope of the adjoint representation $\mathrm{SL}_4 \to \mathrm{GL}(\mathfrak{sl}_4)$.

$B \subset G$ containing T. The proofs of these well-known facts and much more are recorded in [34] (see Section 7). The *fixed subtorus* T_0, and the *isotropic subtorus* T_1 associated with T are defined by

$$T_0 = \{t \in T : \ \theta(t) = t\},$$
$$T_1 = \{t \in T : \ \theta(t) = t^{-1}\}.$$

The multiplication map $T_1 \times T_0 \to T$ is an *isogeny*, that is to say a surjective homomorphism with a finite kernel.

Among all θ-stable maximal tori, we work with the one for which the dimension $l = \dim T_1$ of the isotropic subtorus is maximal. The integer l is called the rank of the symmetric variety G/G^θ. Let Φ denote the set of roots (weights of the adjoint representation) of G relative to T. Passing to the Lie algebra setting by differentiation, we view Φ as a subset of the dual vector space $\mathrm{Lie}(T)^*$ of the Lie algebra of T. Since θ is an automorphism of T, it induces a linear map

$$\theta^* : \ \mathrm{Lie}(T)^* \to \mathrm{Lie}(T)^*,$$

which, in turn induces an involution on Φ. Define

$$\Phi_0 = \{\alpha \in \Phi : \ \theta^*(\alpha) = \alpha\},$$
$$\Phi_1 = \Phi - \Phi_0.$$

Lemma 8 (Lemma 1.2, [7])**.** There exists a system of positive roots $\Phi^+ \subseteq \Phi$ such that $\theta^*(\alpha) \in \Phi - \Phi^+$ for all $\alpha \in \Phi^+ \cap \Phi_1$.

We fix a set of positive roots Φ^+ as in Lemma 8. Let Δ denote the associated set of simple roots and set

$$\Delta_0 = \Phi_0 \cap \Delta,$$
$$\Delta_1 = \Phi_1 \cap \Delta.$$

Observe that $|\Delta_1| \geq \dim T_1 = l$.

It turns out there exists an ordering $\{\alpha_1, \ldots, \alpha_j\}$ of the elements of Δ_1 such that the differences $\alpha_i - \theta^*(\alpha_i)$ are mutually distinct for $i = 1, \ldots, l$, and for each $i \in \{l+1, \ldots, j\}$, there exists an index $s \in [l]$ such that $\alpha_i - \theta^*(\alpha_i) = \alpha_s - \theta^*(\alpha_s)$. A *restricted simple root* $\overline{\alpha}$ is a weight of the form

$$\overline{\alpha} = \frac{\alpha_i - \theta^*(\alpha_i)}{2} \text{ for some } i \in [l].$$

In this case, we denote $\overline{\alpha}$ by $\overline{\alpha}_i$, and denote by $\overline{\Delta_1} = \{\overline{\alpha}_1, \ldots, \overline{\alpha}_l\}$ the set of all restricted simple roots.

Suppose that $\Delta_0 = \{\beta_1, \ldots, \beta_k\}$. In accordance with the partitioning $\Delta = \Delta_0 \sqcup \Delta_1$, we divide the set of fundamental weights of Δ into two disjoint subsets $\{\omega_1, \ldots, \omega_j\} \sqcup \{\zeta_1, \ldots, \zeta_k\}$ so that for each $i \in [j]$ the following holds:

$$(\omega_i, \beta_s^\vee) = 0 \text{ for } s = 1, \ldots, k, \text{ and } (\omega_i, \alpha_r^\vee) = \delta_{i,r} \text{ for } r = 1, \ldots, j.$$

Similarly for ζ_i's. As it is shown in [7] (Page 5 and 6), θ^* induces an involution $\widetilde{\theta}$ on the indices $\{1, \ldots, j\}$ such that $\theta^*(\omega_i) = -\omega_{\widetilde{\theta}(i)}$. Thus, we arrive at a crucial definition for our purposes:

Definition 9. A dominant weight λ of G is called special (or, θ-special), if $\theta^*(\lambda) = -\lambda$. If (ρ, V) is an irreducible representation with a θ-special highest weight, then we call ρ a θ-special representation of G.

Let G_0 be a semisimple, adjoint type linear algebraic group, and let $\theta_0 : G_0 \to G_0$ be an involution on G_0. We choose a θ_0-stable maximal torus T_0 in G_0. Let λ be a special, dominant weight with the corresponding irreducible representation (ρ_0, V). Assume also that (ρ_0, V) is faithful. As before, we define the reductive group G by setting $G = \mathbf{k}^* \cdot \rho_0(G_0) \subset \mathrm{GL}(V)$. We claim that there exists an "extension" of θ_0 to G. To this end we define

$$\theta(c\rho_0(g)) = c^{-1}\rho_0(\theta_0(g)), \qquad g \in G_0,\ c \in \mathbf{k}^*. \tag{3.1}$$

To prove that θ is well defined, suppose $g, g' \in G_0$ and $c, c' \in \mathbf{k}^*$ are such that $c\rho_0(g) = c'\rho_0(g')$. Let $\alpha \in \mathbf{k}^*$ denote cc'^{-1}. Then $\rho_0(g^{-1}g') = \alpha 1_{\mathrm{GL}(V)} \in \mathrm{GL}(V)$. Since V is irreducible, by Schur's Lemma, $g^{-1}g'$ lies in the center of G_0. In other words, $g = g'$, and therefore $c = c'$. Finally, note that

$$\begin{aligned}\theta(\theta(c\rho_0(g))) &= \theta(c^{-1}\rho_0(\theta_0(g))) \\ &= c\rho_0(\theta_0(\theta_0(g))) \\ &= c\rho_0(g) \ \text{ for all } c \in \mathbf{k}^* \text{ and } g \in G_0.\end{aligned}$$

The *antiinvolution* corresponding to θ is, by definition, the composition $\theta_{an} := \theta \circ \iota$ of θ with the "inverting" morphism $\iota : \ g \mapsto g^{-1}$. The map induced by θ_{an} on the weights $X(T)$ is denoted θ^*_{an}. Then θ^* and θ^*_{an} are related to each other by

$$\theta^*_{an}(\chi) = -\theta^*(\chi) \ \text{ for } \chi \in X(T).$$

In particular, if $\theta^*(\lambda) = -\lambda$, then $\theta^*_{an}(\lambda) = \lambda$.

We are ready to prove Theorem 1. Let us paraphrase it for completeness: If M is a J-irreducible monoid that is obtained from a θ-special representation of G, then there exists a unique morphism $\theta_{an} : M \to M$ such that

1. $\theta_{an}(xy) = \theta_{an}(y)\theta_{an}(x)$ for all $x, y \in M$;
2. θ^2_{an} is the identity map on M;
3. $\theta_{an}(g) = \theta(g)^{-1}$ for all $g \in G$.

Proof of Theorem 1. Since θ_{an} agrees with θ (after composing with ι, of course) on G, the uniqueness is clear. We are going to show that θ_{an} extends to whole J-irreducible monoid M associated to an irreducible representation (ρ_0, V) of G_0 with the highest weight λ. Let ρ denote the representation of G as defined in (2.4) (with $k = 1$).

Let T_0 denote the maximal torus of G_0 such that $T = \mathbf{k}^* \cdot \rho_0(T_0)$, and let $\langle \Pi(\rho) \rangle$ denote the submonoid of $X(T)$ generated by the weights $\Pi(\rho)$ of T. The coordinate ring of the affine torus embedding $\overline{T}$ is equal to the monoid-ring $R = \mathbf{k}[\langle \Pi(\rho) \rangle]$ (see [Lemma 3.2, [24]]). Therefore, $\overline{T} = \mathrm{Spec}(R)$. On the other hand, by Lemma 5, we know that $\Pi(\rho)$ is contained in the convex hull $\mathcal{P}$ of $W \cdot (\lambda + \chi)$, where χ is the n-th root of the determinant on $\mathrm{GL}(V)$.

Since λ is special, by [Lemma 1.6, [7]], there is a G-isomorphism $V^\theta \simeq V^*$, hence $\theta^*(\Pi(\rho)) = \Pi(\rho^*) = -\Pi(\rho)$ and it follows $\theta^*_{an}(\Pi(\rho)) = \Pi(\rho)$. In particular, it induces an antiinvolution θ_{an} on $\overline{T} = \mathrm{Spec}(R)$. Since $\theta \circ \iota = \theta_{an}$ on T, by the "extension principle" (see [Corollary 4.5, [24]]), there exists a unique morphism $\theta_{an} : M \to M$, which agrees with $\theta \circ \iota$ on G, and agrees with θ_{an} on $\overline{T}$. Since $\theta^2_{an} = id$ on G, and since G is dense on M, we see that $\theta^2_{an} = id$ on M. Finally, since $(x, y) \mapsto \theta_{an}(xy)$ and $(x, y) \mapsto \theta_{an}(y)\theta_{an}(x)$ are morphisms from $M \times M$ into M agreeing on an open dense set $G \times G$, they agree everywhere.

□

In general, it is important to determine all special dominant weights of a given group. Before presenting some known examples of groups along with the lists of their special weights, we recollect some general information from the representation theory side of the story.

Recall that a closed subgroup $H \subseteq G$ is called spherical, if H has a dense orbit in G/B, or equivalently, B has a dense orbit in G/H. Let $\mathbf{k}[G]$ denote the ring

of regular functions on G. Now, given a spherical subgroup H and an irreducible representation (π, V) of G, the space of H-invariant vectors in V, denoted by V^H, is nonzero if and only if $\dim V^H \leq 1$. Assuming H is reductive, the spherical irreducible representations of G, namely those irreducible representations with $\dim V^H = 1$ are precisely the irreducible representations that occur in

$$\mathbf{k}[G/H] = \{f \in \mathbf{k}[G] : \text{ such that } f(gh) = f(g) \text{ for all } h \in H \text{ and } g \in G\}.$$

Indeed, this follows from the algebraic form of Peter-Weyl theorem: $\mathbf{k}[G/H] = \mathbf{k}[G]^H \simeq \bigoplus V^* \otimes V^H$, where the sum is over all irreducible representations of G. (We thank the referee for pointing it out to us.)

Remark 10. Let V_λ and V_μ be two irreducible representations with highest weights λ and μ, respectively. Let $v \in V_\lambda$ and $w \in V_\mu$ be the corresponding highest weight vectors. If v and w are H-invariant, that is to say if V_λ and V_μ are spherical representations, then $v \otimes w \in V_\lambda \otimes V_\mu$ is H-invariant also. Since $v \otimes w$ is the highest weight vector of the irreducible representation with highest weight $\lambda + \mu$, we see that the sum of two spherical weights is spherical. Hence, spherical dominant weights form a subsemigroup of all dominant weights.

Let λ be a dominant weight and let V_λ denote the corresponding irreducible representation of G with highest weight λ. Let $H \subset G$ denote the fixed subgroup of θ, hence H is spherical. It is shown in Proposition 1.7 of [7] that if λ is special, then the irreducible representation $V_{2\lambda}$ of G with highest weight 2λ has an H-invariant vector, hence 2λ is a spherical weight as well. In fact, the answer to the question "Which spherical dominant weights are special?" is known for quite sometime. The spherical weights coincide with the special weights if and only if G_0/G_0^θ is not of exceptional type. For the complete determination of these weights see page 10 of [7] (or, see Theorem 3 of [37]).

By definition, a *classical group* is one of the following semisimple groups:

- $\mathrm{SL}_n = \{g \in \mathrm{GL}_n : \det g = 1\}$,
- $\mathrm{SO}_n = \{g \in \mathrm{SL}_n : gg^\top = 1\}$,
- $\mathrm{Sp}_{2n} = \{g \in \mathrm{SL}_{2n} : J_n(g^\top)^{-1}J_n = g\}$, where J_n is the $2n \times 2n$ block diagonal matrix $J_n = \mathrm{diag}(J_2, \ldots, J_2)$ with $J_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

There is a well known classification, due to Cartan, of involutions on semisimple groups. For the classical groups, up to inner automorphisms there are seven types of involutions in total. For the exceptional groups there are 9 involutions. See Chapter X, Section 6 of [9] for a complete list. Here we focus only on the classical groups. Following the notation of [8], in Table 1, we list the "classical" symmetric varieties along with the generators of their semigroups of spherical weights.

Classical symmetric variety	Spherical fundamental dominant weights
$\mathrm{SL}_n/\mathrm{SO}_n$	$2\omega_1, 2\omega_2, \ldots, 2\omega_{n-1}$
$\mathrm{SL}_{2n}/\mathrm{Sp}_{2n}$	$\omega_2, \omega_4, \ldots, \omega_{2n-2}$
$\mathrm{SL}_n/S(\mathrm{GL}_p \times \mathrm{GL}_q)$	$\omega_1 + \omega_{n-1}, \omega_2 + \omega_{n-2}, \ldots, \omega_p + \omega_{n-p}$ $(2l \leq n)$
$\mathrm{Sp}_{2n}/\mathrm{GL}_n$	$2\omega_1, 2\omega_2, \ldots, 2\omega_n$
$\mathrm{SO}_{2n}/\mathrm{GL}_n$ (n even)	$\omega_2, \omega_4, \ldots, \omega_{n-2}, 2\omega_n$
$\mathrm{SO}_{2n}/\mathrm{GL}_n$ (n odd)	$\omega_2, \omega_4, \ldots, \omega_{n-3}, \omega_{n-1} + \omega_n$
$\mathrm{SO}_n/S(\mathrm{O}_p \times \mathrm{O}_q)$ $(n = 2k$ even$)$	$2\omega_1, 2\omega_2, \ldots, 2\omega_l$ $(l < k)$ $2\omega_1, 2\omega_2, \ldots, 2\omega_{l-1}, 4\omega_l$ $(l = k)$
$\mathrm{SO}_n/S(\mathrm{O}_p \times \mathrm{O}_q)$ $(n = 2k+1$ odd$)$	$2\omega_1, 2\omega_2, \ldots, 2\omega_l$ $(l < k-1)$ $2\omega_1, 2\omega_2, \ldots, 2\omega_{k-2}, 2\omega_{k-1} + 2\omega_k$ $(l = k-1)$ $2\omega_1, 2\omega_2, \ldots, 2\omega_{k-2}, 2\omega_{k-1}, 2\omega_k$ $(l = k)$
$\mathrm{Sp}_{2n}/(\mathrm{Sp}_{2p} \times \mathrm{Sp}_{2q})$	$2\omega_1, 2\omega_2, \ldots, 2\omega_{2l}$ $(2l \leq n)$

Table 1: Spherical weights for classical symmetric varieties

4 Special compactifications of G_0/G_0^θ

To motivate our next discussion, we focus on the simplest spherical representation for $G_0/G_0^\theta = \mathrm{PSL}_n/\mathrm{PSO}_n$, namely, the irreducible representation of $G_0 = \mathrm{PSL}_n$ corresponding to the doubled first fundamental dominant weight, $2\omega_1$. In this case, the involution on G_0 is given by $\theta(g) = (g^{-1})^\top$. Obviously, the maximal torus of diagonal matrices is θ-stable. Since the action of θ on coordinate functions is given by

$$(\theta \cdot \varepsilon_i)(x) = \varepsilon_i(\theta(x)) = \varepsilon_i(x^{-1}) = t_i^{-1} = \varepsilon_i^{-1}(x),$$

where x is the diagonal matrix $x = \mathrm{diag}(t_1, \ldots, t_n)$, we see that θ^* maps the character ε_i to $-\varepsilon_i$. Here, we are using the additive notation. In particular, since the fundamental dominant weights of (G_0, T_0) are of the form $\omega_i = \sum_{s=1}^{i} \varepsilon_s - \frac{i}{n} \sum_{r=1}^{n} \varepsilon_r$, we see that any element of $X(T_0)$ is special:

$$\theta^* \left(\sum_i^{n-1} a_i \omega_i \right) = - \left(\sum_i^{n-1} a_i \omega_i \right).$$

If V_{ω_i} denotes the irreducible representation with highest weight ω_i, then the homogeneous polynomials of degree k on V_{ω_i} forms an irreducible representation of

G_0, and its T_0-highest weight is $k\omega_i$. In particular, $V_{\omega_1} \cong \mathbf{k}^n$ is irreducible and the highest weight of the second symmetric power $S^2\mathbb{C}^n$ is $2\omega_1$. Concretely, this representation of G_0 comes from the representation of SL_n on the space of $n \times n$ symmetric matrices, which we denote by Sym_n. The action of SL_n on Sym_n is given by

$$g \cdot A = (g^{-1})^\top A g^{-1}, \quad g \in \mathrm{SL}_n, \ A \in \mathrm{Sym}_n. \tag{4.1}$$

Note that an element $A \in \mathrm{Sym}_n$ is fixed by $H = \mathrm{PSO}_n$ if and only if it is (a scalar multiple of) the identity transformation, and that the PSL_n-orbit of the identity consists of matrices of the form $gg^\top$, $g \in \mathrm{PSL}_n$.

For this paragraph only, we specialize our underlying field to $\mathbb{C}$, complex numbers. A well-known result in matrix analysis known as Takagi's factorization states that if $A \in \mathrm{Mat}_n$ is symmetric, then there exists a unitary matrix $U \in \mathrm{Mat}_n$ and a real nonnegative diagonal matrix Σ such that $A = U\Sigma U^\top$. See [12]. In particular, denoting by $\Sigma^{1/2}$ the matrix obtained from Σ by taking the square root of its entries, we see that $A = (U\Sigma^{1/2})(U\Sigma^{1/2})^\top$. In conclusion, any symmetric matrix A has a decomposition of the form $A = BB^\top$ for some $B \in \mathrm{Mat}_n$. Since $\mathrm{GL}_n \simeq \mathbb{C}^*\mathrm{SL}_n$ is dense in Mat_n, we see that the set of all symmetric matrices Sym_n is the closure of the set $\{cgg^\top : \ g \in \mathrm{SL}_n, c \in \mathbb{C}^*\}$. Since $\mathcal{O} = \{gg^\top : \ g \in \mathrm{SL}_n\}$ is equal to $\mathrm{SL}_n \cdot 1_{\mathrm{Mat}_n}$ with respect to the action (4.1), we arrive at the following conclusion:

Sym_n is the embedding of the symmetric variety $\mathbf{k}^* \cdot \mathrm{PSL}_n/\mathrm{PSO}_n \simeq \{cgg^\top : \ g \in \mathrm{PSL}_n, \ c \in \mathbf{k}^*\}$.

The main purpose of this section is to generalize the above observation to all symmetric varieties. We start with recalling a useful construction originally due to De Concini and Procesi.

Lemma 11 ([7], Lemma 1.6)**.** Let λ be a θ-special weight for (G, T), $V = V_\lambda$ be the corresponding irreducible representation and denote by V^θ a copy of V on which G acts by $g \cdot v = \theta(g)v$. Then there is a unique (G,θ)-equivariant isomorphism

$$h : V^* \longrightarrow V^\theta. \tag{4.2}$$

Indeed, normalizing it if necessary, the isomorphism h maps v^λ, a lowest weight vector in the dual space V^* to v_λ, a highest weight vector of V.

Let h, V, v_λ and $v^\lambda \in V^*$ be as in Lemma 11. Extend the singleton $\{v_\lambda\}$ to a basis $\mathcal{V} = \{v_\lambda, v_1, \ldots, v_m\}$ for V where each v_i $(i = 1, \ldots, m)$ is a weight vector. Then there is the dual basis $\mathcal{V}' = \{v^\lambda, v^1, \ldots, v^m\}$ with the property that if χ_i is the weight

corresponding to v_i, then $-\chi_i$ is the weight corresponding to v^i, and furthermore, the weight of $h(v^i)$ is the character $-\theta^*(\chi_i)$.

Since V is self-dual by Lemma 11, there is a G-equivariant isomorphism between $\mathrm{End}(V)$ and $V \otimes V$, defined explicitly as follows: A linear map $f \in \mathrm{Hom}(V,V)$ is first mapped to

$$v^\lambda(f(v_\lambda))(v^\lambda \otimes v_\lambda) + \sum_{i,j=1}^{m} v^i(f(v_j))(v^i \otimes v_j) \in V^* \otimes V$$

via the canonical isomorphism $\mathrm{Hom}(V,V) = V^* \otimes V$, and then it is mapped to

$$v^\lambda(f(v_\lambda))(v_\lambda \otimes v_\lambda) + \sum_{i,j=1}^{m} v^i(f(v_j))(w_i \otimes v_j) \in V \otimes V, \quad \text{where } w_i = h(v^i)$$

by the isomorphism $V^* \otimes V \to V \otimes V$ sending $v' \otimes v \in V^* \otimes V$ to $h(v') \otimes v \in V \times V$.

Remark 12. $*$-action of G on itself is defined by $g * g' = gg'\theta(g)^{-1}$ (for all $g, g' \in G$), and on $V \otimes V$ it is defined by $g * (v_1 \otimes v_2) = gv_1 \otimes \theta(g)^{-1}v_2$ (for all $v_1, v_2 \in V$ and $g \in G$). By the equivariant isomorphism $\pi : \mathrm{End}(V) \to V \otimes V$, the identity element of $\mathrm{End}(V)$ is mapped to $h \in V \otimes V$, where

$$1 = 1_{\mathrm{End}(V)} \rightsquigarrow h = v_\lambda \otimes v_\lambda + \sum_{i=1}^{m} w_i \otimes v_i. \tag{4.3}$$

Thus, we identify the J-irreducible monoid $M = \overline{\mathbf{k}^* \cdot \rho_0(G_0)} \subseteq \mathrm{End}(V)$ with its image in $V \otimes V$, and identify the orbit $P = G * 1_G$ with $G * h$.

The following facts are observed in Section 1.7 of [7]:

- The tensor $v_\lambda \otimes v_\lambda$, which is a weight vector with weight 2λ, generates a copy of $V_{2\lambda}$ in $V \otimes V$.
- $w_i \otimes v_i$'s are weight vectors with $\chi_i - \theta^*(\chi_i)$ as their corresponding weights.
- Under the canonical G-equivariant projection $p : V \otimes V \to V_{2\lambda}$ the "identity tensor" h of (4.3) is mapped onto the vector

  $$h' = v_{2\lambda} + \sum z_i \in V_{2\lambda}, \tag{4.4}$$

  where $v_{2\lambda}$ is the highest weight vector of $V_{2\lambda}$, and z_i's are weight vectors having distinct weights of the form $2(\lambda - \sum_{i=1}^{\ell} n_i\overline{\alpha_i})$ with $n_i \in \mathbb{Z}_{\geq 0}$.
- Both of the elements $h \in V \otimes V$ and $h' \in V_{2\lambda}$ are fixed by $H = G^\theta$.

- If, in addition to being a special weight, λ is regular with respect to G/H (see Definition 1.4, [7]), then the wonderful compactification (defined in the introduction section) of G/H is the Zariski closure $\overline{G \cdot [h']}$ in $\mathbb{P}(V_{2\lambda})$. Here $[h']$ is the point corresponding to the line $\mathbf{k} \cdot h'$ in $V_{2\lambda}$.

Recall that $P = \{g\theta(g)^{-1} : \ g \in G\}$ is a closed embedding of G/G^θ in G (see Section 2 of [28], or see [37]). We are going to describe a special compactification of P along the lines of the construction of its wonderful compactification.

Our assumptions are as before; G_0 is an adjoint semisimple group with an involution θ, λ is a θ-special weight, (ρ_0, V) is the faithful irreducible representation of G_0 with highest weight λ, G is the group $G = \mathbf{k}^* \cdot \rho_0(G_0) \subset \mathrm{GL}(V)$ on which the involution θ extends as in (3.1), and finally $P = \{g\theta(g)^{-1} : \ g \in G\}$, $P_0 = \{g\theta(g)^{-1} : \ g \in G_0\}$.

Since P is the orbit of h in $V \otimes V$ and since $p : V \otimes V \to V_{2\lambda}$ is G-equivariant, P is mapped isomorphically onto its image $p(P) = G * h'$ in $V_{2\lambda}$. Note that neither P, nor its image $p(P)$ contain 0. Note also that ρ_0 is assumed to be faithful, so we identify P_0 with its image $p(\rho_0(P_0))$ in $V_{2\lambda}$.

Notation: Let A be a subset of an arbitrary vector space W. We denote by $\mathbb{P}(A; W)$ the image of $A - \{0\}$ in $\mathbb{P}(W)$.

Definition 13. The special compactification X_{P_0} of P_0 is the normalization of the Zariski closure $\overline{\mathbb{P}(p(P); V_{2\lambda})} \subset \mathbb{P}(V_{2\lambda})$.

Example 14. Consider ω_1 of $G_0 = \mathrm{PSL}_n$ with $\theta(g) = (g^{-1})^\top$. Then $G = \mathbf{k}^* \cdot \rho_0(G_0)$ is GL_n and $M = \mathrm{Mat}_n = \mathrm{End}(\mathbf{k}^n)$, and the orbit $G \cdot 1_{\mathrm{Mat}_n}$ is equal to $P = \mathbf{k}^* \cdot \{gg^\top : g \in \mathrm{SL}_n\}$, the set of all $n \times n$ invertible symmetric matrices. Clearly, the closure $\overline{P}$ of this set in M is equal to Sym_n. Since $V_{2\omega_1} \simeq \mathrm{Sym}_n$, the normalization in $\mathbb{P}(V_{2\omega_1})$ of the Zariski closure of the image of P is the projective space itself;

$$\mathbb{P}(\mathrm{Sym}_n) = (\mathrm{Sym}_n - \{0\})/\sim .$$

Here, we write $[A] \sim [B]$ for $A, B \in \mathrm{Sym}_n$, if there exists $c \in \mathbf{k}^*$ such that $A = cB$.

5 Borel orbits in special compactifications

In the rest of our paper we describe the parametrizing set of the orbits of a Borel subgroup in a special compactification. We start with gathering some useful facts.

On one hand, the internal structure of a Borel subgroup $B \subseteq G$ is very simple; it is the semi-direct product of a maximal torus T and a maximal unipotent subgroup U, which is normalized by T. On the other hand, the role played by B within G is huge; when G is decomposed into double cosets of B the parametrizing set is the Weyl group $W = N_G(T)/T$. Recall (from Section 2.2) that there is an extension of

this decomposition to the reductive monoids. Since we need the notation from these results, we elaborate:

Lemma 15 (Generalized Bruhat-Chevalley decomposition, [25])**.** Let M be a reductive monoid with the group of invertible elements G, and let $T \subseteq B$ be a maximal torus contained in a Borel subgroup. Let $\overline{N}$ denote the closure in M of the normalizer $N = N_G(T)$ of T. If $m \in M$, then there exist $b_1, b_2 \in B$, and $\overline{n} \in \overline{N}$ such that $m = b_1 \overline{n} b_2$. This leads to the Bruhat-Chevalley decomposition of M:

$$M = \bigcup_{\overline{n} \in R} B\overline{n}B, \tag{5.1}$$

where the union is disjoint and $R = \overline{N}/T$ is the Renner monoid of M.

Fix an element $\overline{n} \in \overline{N}$, and let $V_{\overline{n}} \subseteq U$ denote the subgroup $V_{\overline{n}} = \{u \in U : u\overline{n}B \subseteq \overline{n}B\}$. Notice that T is contained the normalizer of $V_{\overline{n}}$. Since $V_{\overline{n}}$ is a closed subgroup of U, there exists a complementary[2] subgroup

$$U_{\overline{n},1} = \prod_{U_\alpha \not\subseteq V_{\overline{n}}} U_\alpha. \tag{5.2}$$

Similarly, let $Z_{\overline{n}} \subseteq U$ denote the closed subgroup $Z_{\overline{n}} = \{u \in U : \overline{n}Tu = \overline{n}T\}$. Also in this case, T is contained in the normalizer of $Z_{\overline{n}}$. Let

$$U_{\overline{n},2} = \prod_{U_\alpha \not\subseteq Z_{\overline{n}}} U_\alpha \tag{5.3}$$

denote its complementary subgroup. The precise structure of the orbit $B\overline{n}B$ is exhibited in the next result:

Lemma 16 (Lemma 13.1, [27])**.** The product morphism $U_{\overline{n},1} \times \overline{n}T \times U_{\overline{n},2} \to B\overline{n}B$ is an isomorphism of varieties.

As a consequence of Lemmas 15 and 16 we have the following important observation:

Uniqueness Criterion: Given an element $m \in M$, there exist unique $u \in U_{\overline{n},1}, v \in U_{\overline{n},2}$, and $\overline{n} \in \overline{N}$ such that

$$m = u\overline{n}v. \tag{5.4}$$

[2]Complementary in this context means that the product morphism $U_{\overline{n},1} \times V_{\overline{n}} \to U$ is an isomorphism of algebraic groups

We continue with the assumption that M is a reductive monoid with an antiinvolution $\theta_{an} : M \to M$. Let $H \subseteq G$ denote, as usual, the fixed subgroup G^θ, where $\theta : G \to G$ is the involution $\iota \circ \theta_{an}$, where ι stands for the inverse map. Recall that the $*$-action of G on M is defined in (1.2). Here, we are going to investigate the sets of $B*$-orbits in following varieties:

- the Zariski closure $\overline{P}$ in M of $P = \{g\theta(g^{-1}) : \ g \in G\} \simeq G/H$;
- the Zariski closure $\overline{Q}$ in M of $Q = \{g \in G : \ \theta(g) = g^{-1}\}$;
- and $M_Q := \{x \in M : \ \theta_{an}(x) = x\}$.

Here, we should mention that the $B*$-orbits in $\overline{P}$ correspond to the B-orbits in $\overline{\mathbb{P}(p(P); V_{2\lambda})} \subset \mathbb{P}(V_{2\lambda})$ (see Definition 13).

Assume from now on that T is a θ-stable maximal torus of the θ-stable Borel subgroup $B \subseteq G$. Notice in this case that the corresponding unipotent subgroup $U \subset B$ has to be θ-stable, as well. Moreover, since T is θ-stable, if n is an element from the normalizer $N = N_G(T)$, then $\theta(n)t\theta(n)^{-1} = \theta(n)\theta(t')\theta(n^{-1}) = \theta(nt'n^{-1}) \in T$ for some $t' \in T$. In other words, $\theta(N) = N$. It follows that the Zariski closure $\overline{N}$ is θ_{an}-stable.

Proposition 17. Any $B*$-orbit in M_Q contains an element of $\overline{N}$.

Proof. For $m \in M_Q$, as it is shown in the previous section, there exist unique $u \in U_{\overline{n},1}, v \in U_{\overline{n},2}$, and $\overline{n} \in \overline{N}$ such that $m = u\overline{n}v$. Then

$$u\overline{n}v = m = \theta_{an}(m) = \theta_{an}(v)\theta_{an}(\overline{n})\theta_{an}(u) = \theta(v)^{-1}\theta_{an}(\overline{n})\theta(u)^{-1}.$$

Since Bruhat-Chevalley decomposition (5.1) is a disjoint union, we see that $\theta_{an}(\overline{n}) = \overline{n}$. Let a denote $v\theta(u)$. It is clear that $\overline{n}a$ lies in the $B*$-orbit of m. Therefore, $\overline{n}a = \theta_{an}(\overline{n}a) = \theta_{an}(a)\theta_{an}(\overline{n}) = \theta(a)^{-1}\overline{n}$ implies

$$\theta(a)\overline{n}a = \overline{n}. \tag{5.5}$$

Suppose $\overline{n} = en$ for some $e \in E(\overline{T})$ and $n \in N$. By (5.5) we see that $\theta(a)e = ena^{-1}n^{-1}$. It follows that the element $\theta(a)e = e\theta(a)e = ena^{-1}n^{-1}e = ena^{-1}n^{-1}$ is contained in the unit group of eMe. In particular, $e\theta(a)e$ lies in the unipotent radical of the Borel subgroup eBe of the $\mathscr{H}$-class of e. See Corollary 7.2 (ii) [21]. Since square roots exists in unipotent groups, we see that $(e\theta(a)e)^{1/2} = e\theta(a)^{1/2}e = (ena^{-1}n^{-1}e)^{1/2} = e(na^{-1}n^{-1})^{1/2}e = e(na^{-1/2}n^{-1})e$. Since $eBe = eC_G(e)$, we have that $e\theta(a)^{1/2}e = \theta(a)^{1/2}e$ and that $e(na^{-1/2}n^{-1})e = ena^{-1/2}n^{-1}$. Now, on one hand we have $\theta(a)^{1/2}e = ena^{-1/2}n^{-1}$, or equivalently $\theta(a)^{1/2}\overline{n}a^{1/2} = \overline{n}$. On the other hand, $\theta(a)^{1/2}\overline{n}a^{1/2} = \theta(a)^{1/2} * (\overline{n}a)$. Therefore, $\overline{n}$ is contained in the $B*$-orbit of m. □

There is an obvious extension to M of the map (1.1) of Vust:

$$\tau : M \longrightarrow M$$
$$\tau(x) = x\theta_{an}(x).$$

We claim that τ and θ_{an} commute with each other. Indeed, if $m \in M$, then

$$\tau(\theta_{am}(m)) = \theta_{an}(m)\theta_{an}(\theta_{an}(m)) = \theta_{an}(m\theta_{an}(m)) = \theta_{an}(\tau(m)).$$

Therefore, if $A \subseteq M$ is θ_{an}-stable, then so are $\tau(A)$ and $\tau^{-1}(A)$. In particular, $\tau^{-1}(\overline{N})$ is θ_{an}-stable.

Lemma 18. *$T \times H$ acts on $\tau^{-1}(\overline{N})$ by $(t,h) \cdot m = tmh^{-1}$.*

Proof. It suffices to check that $\tau(tmh^{-1}) = tm\theta_{an}(m)\theta(t)^{-1}$ is in $\overline{N}$ whenever $m \in \tau^{-1}(\overline{N})$. This is straightforward. □

Theorem 19. The following sets are in bijection with each other;

1. $B*$-orbits in M_Q, and
2. $T \times H$-orbits in $\tau^{-1}(\overline{N}) \subset M$.

Proof. We claim that $B \times H$-orbits in M are in one-to-one correspondence with $B*$-orbits in M_Q. Indeed, if $a \in M$, then

$$\tau(bah^{-1}) = bah^{-1}\theta_{an}(bah^{-1}) = ba\theta_{an}(a)\theta_{an}(b) = b * \tau(a).$$

Conversely, for $x \in M_Q$, $\tau(BaH) = B * x$ for any $a \in M$ such that $x = \tau(a)$.

By Proposition 17, we know that each $B*$-orbit intersect $\overline{N}$ non-trivially. Suppose $\overline{n}_1$ and $\overline{n}_2$ are two elements of $\overline{N}$ representing the same $B*$-orbit. Then there exists $b \in B$ such that $\overline{n}_1 = b\overline{n}_2\theta_{an}(b)$. Since the decomposition $M = \bigcup_{\dot{r} \in R} BrB$ is a disjoint union, $b = t \in T$ for some $t \in T$. Moreover, for any $t \in T$, $B * (t\overline{n}\theta_{an}(t)) = B * \overline{n}$. Therefore, $B*$-orbits in M_Q are parametrized by the $T*$-orbits in $\overline{N} \cap M_Q$. But these orbits are in one-to-one correspondence with $T \times H$ orbits in $\tau^{-1}(\overline{N})$.

□

Now we are ready to prove our second main result, which states that the following sets are finite and they are in bijection with each other:

1. $B*$-orbits in $\overline{Q}$ (respectively, $B*$-orbits in $\overline{P}$),
2. $T \times H$-orbits in $\tau^{-1}(\overline{N} \cap \overline{Q})$ (respectively, $T \times H$-orbits in $\tau^{-1}(\overline{N} \cap \overline{P})$).

Proof of Theorem 2. Finiteness follows from the fact that the sets $\overline{P}$ and $\overline{Q}$ are spherical varieties. The rest is immediate from Theorem 19 and the fact that $P \subseteq Q \subseteq M_Q$ are $B*$-stable. □

We finish this section by presenting two important examples.

Example 20. Going back to Example 14, we see that $Q = \{g \in \mathrm{PSL}_n : \ g = g^\top\}$ of $\mathrm{PSL}_n/\mathrm{PSO}_n$ coincides with $P = \{gg^\top : \ g \in \mathrm{PSL}_n\}$, and therefore, $\overline{P} = \overline{Q}$ in $M = \mathrm{Mat}_n$ is equal to Sym_n. Also, the unique (anti)-extension $\theta_{an} : \mathrm{Mat}_n \to \mathrm{Mat}_n$ of $\theta(g) = (g^\top)^{-1}$, $g \in \mathrm{GL}_n$ is $\theta_{an}(x) = x^\top$, $x \in \mathrm{Mat}_n$. Therefore, M_Q is equal to $\mathrm{Sym}_n \subset \mathrm{Mat}_n$, also. Finally, we know from [35] that $B*$-orbits in Sym_n are parameterized by $n \times n$ *partial involutions*, that is to say, symmetric $0/1$ matrices with at most one 1 in each row and each column.

Example 21. Let $V = \bigwedge^2 \mathbf{k}^{2n}$ denote the second exterior power of $\mathbf{k}^{2n}$. Then, according to Table 1, (ρ_0, V) is the irreducible representation of $G_0 = \mathrm{PSL}_{2n}$ with special T_0-highest weight ω_2. Here, T_0 is the maximal torus consisting of diagonal matrices in G_0, and the involution is $\theta'(g) = -J(g^{-1})^\top J$, where J is the $2n \times 2n$ block diagonal matrix

$$J = \mathrm{diag}(J_2, \ldots, J_2) \ \text{ with } J_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

Explicitly V is the space of $2n \times 2n$ skew-symmetric matrices, and the action of G_0 on V is

$$g \cdot A = (g^{-1})^\top A g^{-1}.$$

For notational ease, let us denote the operator $\rho_0(g) \in \mathrm{GL}(V)$, $g \in G_0$ by ϕ_g. Note that $\phi_J^2 = 1_{\mathrm{GL}(V)}$. It is not difficult to show that ρ_0 is faithful, and that the extension of θ' to $G = \mathbf{k}^* \cdot \rho_0(G_0)$ is given by $\theta(c\phi_g) = c\phi_J \phi_{(g^{-1})^\top} \phi_J$, where $c \in \mathbf{k}^*$. Next we are going to take a closer look at the (closures of the) varieties $P = \{y\theta(y)^{-1} : \ y \in G\}$ and $Q = \{y \in G : \ y^{-1} = \theta(y)\}$.

If $y = c\phi_g \in Q$ for some $g \in G_0$, and $c \in \mathbf{k}^*$, then

$$c^{-1}\phi_g^{-1} = \theta(y) = c\phi_J \phi_{(g^{-1})^\top} \phi_J, \quad \text{or, equivalently} \quad 1_V = c^2 \phi_{gJ(g^{-1})^\top J}.$$

Since ρ_0 is faithful, $c = 1$, and $gJ(g^{-1})^\top J = 1_{G_0}$, or $g^{-1} = J(g^{-1})^\top J$. In other words, Q is isomorphic to $Q_0 = \{g \in \mathrm{PSL}_{2n} : \ \theta'(g) = g^{-1}\}$. On the other hand, it is known that Q_0 is equal to $P_0 = \{g\theta'(g^{-1}) : \ g \in \mathrm{PSL}_{2n}\}$. See Section 11.3.5 of [8]. Since the image of P_0 under ρ_0 is equal to P, we see that $P = Q$, and therefore, P is a closed subvariety of G. Let θ_{an} be the unique antiinvolution extension of θ to the monoid M of (ρ_0, V). Then the closure M_Q of Q in M is equal to $M_Q = \{y \in M : \ \theta_{an}(y) = y\}$.

Next, we compute the parametrizing set of $B*$-orbits in $M_Q = M_P$. To this end, we determine the normalizer of T in G. We claim that $N_G(T) = \mathbf{k}^* \rho_0(N_{G_0}(T_0))$. Indeed, let $x = c\rho_0(g) \in G$ be an element from the normalizer of T, and let $t \in T$. Since $t = d\rho_0(t')$ for some $t' \in T_0$ and $d \in \mathbf{k}^*$, we have $xtx^{-1} = d\rho_0(gt'g^{-1}) \in T$, or equivalently, $gt'g^{-1} \in T_0$. Thus, t lies in $\mathbf{k}^*\rho_0(N_{G_0}(T_0))$. The other inclusion is obvious. We continue with analyzing a typical element of $N_G(T)$. Assume that g is a monomial matrix, that is to say, every row and every column have exactly one nonzero entry. We are going to prove that, once we fix a basis, $\rho_0(g) = \phi_g$ is a monomial matrix, also. For this purpose, we choose the following basis

$$F_{i,j} = E_{i,j} - E_{j,i} \qquad 1 \leq i < j \leq 2n,$$

where $E_{i,j}$'s are the elementary matrices. Suppose that the inverse of $g \in G_0$ is the matrix $g^{-1} = (g_{k,l})_{k,l=1}^{2n}$. Obviously, g^{-1} is a monomial matrix, as well. Since $\rho_0(g) \cdot E_{i,j} = (g^{-1})^\top E_{i,j} g^{-1} = (g_{i,k}g_{j,l})_{k,l}^{2n}$, we see that

$$g \cdot F_{i,j} = \rho_0(g) \cdot F_{i,j} = (g_{i,k}g_{j,l} - g_{j,k}g_{i,l})_{k,l}^{2n}. \tag{5.6}$$

We continue with a special case of our claim by assuming that g is a diagonal matrix. Then the k,l-th entry (with $k < l$) of $g \cdot F_{i,j}$ is nonzero if and only if $i = k$ and $j = l$. In this case, $g \cdot F_{i,j} = g_{i,i}g_{j,j}F_{i,j}$. Thus, the matrix representing $\rho_0(g)$ is the $n(n-1) \times n(n-1)$ diagonal matrix $\mathrm{diag}(s_{1,2}, s_{1,3}, \ldots, s_{n-1,n})$ with $s_{i,j} = g_{i,i}g_{j,j}$. Now, more generally, assume that g is a monomial matrix. Then the k,l-th entry (with $k < l$) $g_{i,k}g_{j,l} - g_{j,k}g_{i,l}$ of $g \cdot F_{i,j}$ is nonzero if and only if one of the following is true;

i) the entries of g^{-1} at its i,k-th and the j,l-th positions are nonzero at the same time, or

ii) the entries of g^{-1} at its i,l-th and the j,k-th positions are nonzero at the same time.

Observe that i) and ii) do not hold true at the same time. Observe also that, for each $i < j$ there exists a unique pair (k,l) with $k < l$ such that either i) is true, or ii) is true. Therefore, if g^{-1} is a monomial matrix, then

$$g \cdot F_{i,j} = \begin{cases} g_{i,k}g_{j,l}F_{k,l} & \text{if } g_{i,k}g_{j,l} \neq 0, \\ g_{i,l}g_{j,k}F_{k,l} & \text{if } g_{i,l}g_{j,k} \neq 0. \end{cases} \tag{5.7}$$

It follows that if g is a monomial matrix, then so is the matrix of $\rho_0(g) = \phi_g$.

Now, let $x \in M$ be an element from $\overline{N_G(T)}$. Since the elements of $\overline{N_G(T)}$ are obtained from those of $N_G(T)$ by taking limits (in the algebraic sense), we see $x \cdot F_{i,j}$

is either identically zero, or it is a scalar multiple of $F_{k,l}$ for some k, l as in (5.7). In other words, x is obtained from the image of a monomial matrix in G_0 by replacing some of its entries by zeros.

It is well know that the invertible symmetric monomial matrices modulo the maximal torus of diagonal matrices represent the $B*$-orbits in Q, and furthermore, the finite set of orbit representatives is in bijection with the fixed point free involutions of the symmetric group S_{2n} (see [29]). Thus, in our case, the representing matrices are those that are obtained from the fixed point free monomial matrices by replacing some of the nonzero entries by zeros. These are precisely the "partial fixed point free involutions," introduced in [6]. See also [4].

6 Final remarks

Given a reductive monoid M with an antiinvolution θ_{an}, we now have the notion of a *symmetric submonoid*

$$M^{an} := \{m \in M : m\theta_{an}(m) = 1_M\}. \tag{6.1}$$

Observe that the identity element 1_M of M is the identity element 1_G of G. Therefore, $\theta_{an}(1_M) = \theta_{an}(1_G) = \theta_{an}(1_G)\theta_{an}(1_G)$, hence $\theta_{an}(1_M) = 1_M$. In other words, $1_M \in M^{an}$. Also, if $m_1, m_2 \in M^{an}$, then

$$m_1m_2\theta_{an}(m_1m_2) = m_1m_2\theta_{an}(m_2)\theta_{an}(m_1) = m_1 \cdot 1_M \cdot \theta_{an}(m_1) = 1_M.$$

Therefore, $m_1m_2 \in M^{an}$. Note that if an element $g \in G$ lies in M^{an}, then $1_G = g^{-1}\theta_{an}(g^{-1}) = g^{-1}\theta(g)$, hence $\theta(g) = g$. In other words, the group of invertible elements of M^{an} is the fixed subgroup $H = G^\theta$. The above argument provides us with an effective way of producing new linear algebraic monoids, namely, one for each antiinvolution θ_{an} on M.

References

[1] Michel Brion. Quelques propriétés des espaces homogènes sphériques. *Manuscripta Math.*, 55(2):191–198, 1986.

[2] Michel Brion. On orbit closures of spherical subgroups in flag varities. *Comment. Math. Helv.*, 76(2):263–299, 2001.

[3] Michel Brion. On algebraic semigroups and monoids. In *Algebraic Monoids, Group Embeddings and Algebraic Combinatorics*, volume 71 of *Fields Institute Communication Series*, pages 1–55. Springer, 2014.

[4] Mahir Bilen Can, Yonah Cherniavsky, and Tim Twelbeck. Bruhat order on partial fixed point free involutions. *Electron. J. Combin.*, 21(4):Paper 4.34, 23, 2014.

[5] Élie Cartan. Sur une classe remarquable d'espaces de Riemann. II. *Bull. Soc. Math. France*, 55:114–134, 1927.

[6] Yonah Cherniavsky. On involutions of the symmetric group and congruence B-orbits of anti-symmetric matrices. *Internat. J. Algebra Comput.*, 21(5):841–856, 2011.

[7] Corrado De Concini and Claudio Procesi. Complete symmetric varieties. In *Invariant theory (Montecatini, 1982)*, volume 996 of *Lecture Notes in Math.*, pages 1–44. Springer, Berlin, 1983.

[8] Roe Goodman and Nolan R. Wallach. *Symmetry, representations, and invariants*, volume 255 of *Graduate Texts in Mathematics*. Springer, Dordrecht, 2009.

[9] Sigurdur Helgason. *Differential geometry, Lie groups, and symmetric spaces*, volume 34 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001. Corrected reprint of the 1978 original.

[10] Aloysius G. Helminck. Computing B-orbits on G/H. *J. Symbolic Comput.*, 21(2):169–209, 1996.

[11] Aloysius G. Helminck and Shu Ping Wang. On rationality properties of involutions of reductive groups. *Adv. Math.*, 99(1):26–96, 1993.

[12] Roger A. Horn and Charles R. Johnson. *Matrix analysis*. Cambridge University Press, Cambridge, 1990. Corrected reprint of the 1985 original.

[13] Yongdong Huang and Naichung Conan Leung. A uniform description of compact symmetric spaces as Grassmannians using the magic square. *Math. Ann.*, 350(1):79–106, 2011.

[14] James E. Humphreys. *Linear algebraic groups*. Springer-Verlag, New York, 1975. Graduate Texts in Mathematics, No. 21.

[15] James E. Humphreys. *Introduction to Lie algebras and representation theory*, volume 9 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1978. Second printing, revised.

[16] Zhenheng Li, Zhuo Li, and You'an Cao. Reductive monoids and renner monoids. In *Algebraic Monoids, Group Embeddings and Algebraic Combinatorics*, volume 71 of *Fields Institute Communication Series*, pages 141–189. Springer, 2014.

[17] Toshihiko Matsuki. The orbits of affine symmetric spaces under the action of minimal parabolic subgroups. *J. Math. Soc. Japan*, 31(2):331–357, 1979.

[18] Toshihiko Matsuki. Closure relations for orbits on affine symmetric spaces under the action of parabolic subgroups. Intersections of associated orbits. *Hiroshima Math. J.*, 18(1):59–67, 1988.

[19] Claus Mokler. Extending the Bruhat order and the length function from the Weyl group to the Weyl monoid. *J. Algebra*, 275(2):815–855, 2004.

[20] Vladimir L. Popov. Contractions of actions of reductive algebraic groups. *Mat. Sb. (N.S.)*, 130(172)(3):310–334, 431, 1986.

[21] Mohan S. Putcha. *Linear algebraic monoids*, volume 133 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1988.

[22] Mohan S. Putcha. Monoid Hecke algebras. *Trans. Amer. Math. Soc.*, 349(9):3517–3534, 1997.

[23] Mohan S. Putcha. Bruhat-Chevalley order in reductive monoids. *J. Algebraic Combin.*, 20(1):34–53, 2004.

[24] Lex E. Renner. Classification of semisimple algebraic monoids. *Trans. Amer. Math. Soc.*, 292(1):193–223, 1985.

[25] Lex E. Renner. Analogue of the Bruhat decomposition for algebraic monoids. *J. Algebra*, 101(2):303–338, 1986.

[26] Lex E. Renner. An explicit cell decomposition of the wonderful compactification of a semisimple algebraic group. *Canad. Math. Bull.*, 46(1):140–148, 2003.

[27] Lex E. Renner. *Linear algebraic monoids*, volume 134 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 2005. Invariant Theory and Algebraic Transformation Groups, V.

[28] Roger W. Richardson. On orbits of algebraic groups and Lie groups. *Bull. Austral. Math. Soc.*, 25(1):1–28, 1982.

[29] Roger W. Richardson and Tonny A. Springer. The Bruhat order on symmetric varieties. *Geom. Dedicata*, 35(1-3):389–436, 1990.

[30] Alvaro Rittatore. Algebraic monoids and group embeddings. *Transform. Groups*, 3(4):375–396, 1998.

[31] Louis Solomon. The Bruhat decomposition, Tits system and Iwahori ring for the monoid of matrices over a finite field. *Geom. Dedicata*, 36(1):15–49, 1990.

[32] Louis Solomon. An introduction to reductive monoids. In *Semigroups, formal languages and groups (York, 1993)*, volume 466 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, pages 295–352. Kluwer Acad. Publ., Dordrecht, 1995.

[33] Tonny A. Springer. Some results on algebraic groups with involutions. In *Algebraic groups and related topics (Kyoto/Nagoya, 1983)*, volume 6 of *Adv. Stud. Pure Math.*, pages 525–543. North-Holland, Amsterdam, 1985.

[34] Robert Steinberg. *Endomorphisms of linear algebraic groups*. Memoirs of the American Mathematical Society, No. 80. American Mathematical Society, Providence, R.I., 1968.

[35] Fernando Szechtman. Equivalence and congruence of matrices under the action of standard parabolic subgroups. *Electron. J. Linear Algebra*, 16:325–333 (electronic), 2007.

[36] Ernest B. Vinberg. Complexity of actions of reductive groups. *Funktsional. Anal. i Prilozhen.*, 20(1):1–13, 96, 1986.

[37] Thierry Vust. Opération de groupes réductifs dans un type de cônes presque homogènes. *Bull. Soc. Math. France*, 102:317–333, 1974.